



(voh-SEV-ee)

## MOST IMPORTANT INFORMATION ABOUT VOSEVI

### **VOSEVI can cause serious side effects, including:**

**Hepatitis B virus reactivation:** Before starting treatment with VOSEVI, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during or after treatment with VOSEVI. This may cause serious liver problems including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B reactivation during and after you stop taking VOSEVI.

## ABOUT VOSEVI

VOSEVI is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C (Hep C) infection with or without cirrhosis (compensated) who have:

- genotype 1, 2, 3, 4, 5, or 6 infection and have previously been treated with a Hep C regimen containing an NS5A inhibitor.
- genotype 1a or 3 infection and have previously been treated with a Hep C regimen containing sofosbuvir without an NS5A inhibitor.

**Do not take VOSEVI** if you take medicines that contain rifampin (Rifater®, Rifamate®, Rimactane®, Rifadin®).

## POSSIBLE SIDE EFFECTS OF VOSEVI

### **VOSEVI can cause serious side effects, including:**

- **Those in the “Most Important Information about VOSEVI” section.**
- **Slow heart rate (bradycardia):** VOSEVI, when taken with amiodarone (Cordarone®, Nexterone®, Pacerone®), a medicine used to treat certain heart problems, may cause serious side effects, including slow heart rate. In some cases slow heart rate has led to death or the need for a pacemaker when amiodarone is taken with medicines containing sofosbuvir. Get medical help right away if you take amiodarone with VOSEVI and get any of the following symptoms:
  - fainting or near-fainting
  - extreme tiredness
  - dizziness or lightheadedness
  - shortness of breath
  - not feeling well
  - chest pains
  - weakness
  - confusion
  - memory problems

**The most common side effects of VOSEVI include** headache, tiredness, diarrhea, and nausea.

These are not all the possible side effects of VOSEVI. Tell your healthcare provider if you have any new symptoms while taking VOSEVI.

## IMPORTANT FACTS

This is only a brief summary of important information about VOSEVI and does not replace talking to your healthcare provider about your condition and your treatment.

## BEFORE TAKING VOSEVI

### **Tell your healthcare provider about all of your medical conditions, including if you:**

- have ever had hepatitis B infection
- have liver problems other than hepatitis C infection
- have severe kidney problems or are on dialysis
- are pregnant or breastfeeding, or plan to become pregnant or breastfeed

### **Tell your healthcare provider about all the medicines you take:**

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider.

VOSEVI and certain medicines may affect each other, or cause side effects.

## HOW TO TAKE VOSEVI

- Take VOSEVI with food.
- If you need to take an antacid that contains aluminum or magnesium, take it 4 hours before or 4 hours after you take VOSEVI.

## GET MORE INFORMATION

- This is only a brief summary of important information about VOSEVI. Talk to your healthcare provider or pharmacist to learn more.
- Go to [VOSEVI.com](http://VOSEVI.com) or call 1-833-4-VOSEVI
- To learn about potential savings, call 1-855-7-MYPATH or go to [VOSEVI.com/support-path](http://VOSEVI.com/support-path)

