



CONVERSATION GUIDE



You're determined to put hepatitis C (Hep C) behind you. VOSEVI is a Hep C treatment specifically for adults who have been previously treated with an advanced Hep C regimen,* but haven't been cured. Cure means the Hep C virus is not detected in the blood when measured three months after treatment is completed.

VOSEVI is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C (Hep C) genotype 1, 2, 3, 4, 5, or 6 infection with or without cirrhosis (compensated) who have previously been treated with a Hep C regimen containing an NS5A inhibitor.

*A Hep C regimen containing an NS5A inhibitor, including EPCLUSA®, HARVONI®, Viekira Pak®, Viekira XR®, Technivie®, Zepatier®, and regimens containing Daklinza®.

Learning if VOSEVI is right for you starts with speaking with your Hep C Specialist.

1

Tell your Hep C Specialist your medical history, including your history with Hep C and any previous Hep C treatment. This will help your Hep C Specialist determine if VOSEVI may be right for you.

Write down your medical history. Include anything that may be affecting your health right now, such as liver problems other than Hep C, kidney problems or if you are on dialysis, or any other medical condition:

2

Tell your Hep C Specialist about all the medicines you're currently taking, including prescription and over-the-counter medicines. Be sure to include antacids, as well as vitamins and herbal supplements. VOSEVI and certain other medicines may affect each other, or may cause side effects.

List all the medicines (prescription and over-the-counter), antacids, vitamins, and herbal supplements you take:

Important Safety Information

What is the most important information I should know about VOSEVI?

VOSEVI can cause serious side effects, including:

- **Hepatitis B virus reactivation:** Before starting VOSEVI treatment, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during and after treatment with VOSEVI. This may cause serious liver problems including liver failure and death. If you are at risk, your healthcare provider will monitor you during and after taking VOSEVI.

Please see Important Facts about VOSEVI including Important Warning on hepatitis B reactivation located on page 4.

3

Share any problems you may have had with staying on treatment in the past. Your Hep C Specialist can work with you to make a plan to help you take your medicine every day.
Write down your concerns about staying on treatment:

4

Keep asking questions. If there's something you don't understand, it's OK to keep asking until you do. Your Hep C Specialist wants you to succeed as much as you do.
Write down any additional questions you have for your Hep C Specialist:

VOSEVI SUPPORT PATH[®] IS ON YOUR SIDE

If you and your Hep C Specialist decide that VOSEVI is right for you, VOSEVI Support Path program navigators are ready to provide information and support to help you get started on treatment.



Connect with
VOSEVI Support Path

1-855-7-MYPATH
(1-855-769-7284)

Important Safety Information (continued)

Who should not take VOSEVI?

- **Do not take VOSEVI** if you take medicines that contain rifampin (Rifater[®], Rifamate[®], Rimactane[®], Rifadin[®]).

What should I tell my healthcare provider before taking VOSEVI?

- **Tell your healthcare provider about all of your medical conditions**, including if you have ever had hepatitis B infection or liver problems other than hepatitis C infection; if you have severe kidney problems or are on dialysis; or if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed. It is not known if VOSEVI will harm your unborn baby or pass into your breast milk.
- **Tell your healthcare provider and pharmacist about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. VOSEVI and certain other medicines may affect each other, or may cause side effects.

Please see Important Facts about VOSEVI including Important Warning on hepatitis B reactivation located on page 4.

With Hep C treatment, the most important thing a person can do for their best chance of achieving a cure is to not miss taking any of their medication—and to finish all of it. Use the tips below to help you stay on track.

TIPS TO HELP YOU REACH YOUR GOAL



Take VOSEVI once a day exactly as prescribed.



Take VOSEVI with food at the same time every day.



Keep VOSEVI with your other daily medications.



Set your alarm clock or phone alarm as a daily reminder.



Leave a reminder note where you'll be sure to see it.

Remember, you haven't been left behind: **Your cure matters.**

To speak with a VOSEVI Educator call **1-833-4-VOSEVI** (1-833-486-7384)

Important Safety Information (continued)

What are the possible side effects of VOSEVI?

Serious side effects may also include:

- **Slow heart rate (bradycardia):** VOSEVI, when taken in combination with amiodarone (Cordarone[®], Nexterone[®], Pacerone[®]), a medicine used to treat certain heart problems, may cause slow heart rate. In some cases slow heart rate has led to death or the need for a pacemaker when amiodarone is taken with medicines containing sofosbuvir. Get medical help right away if you take amiodarone with VOSEVI and get any of the following symptoms: fainting or near-fainting, dizziness or lightheadedness, not feeling well, weakness, extreme tiredness, shortness of breath, chest pains, confusion, or memory problems.

The most common side effects of VOSEVI include headache, tiredness, diarrhea, and nausea.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Facts about VOSEVI including Important Warning on hepatitis B reactivation located on page 4.



(voh-SEV-ee)

MOST IMPORTANT INFORMATION ABOUT VOSEVI

VOSEVI can cause serious side effects, including:

Hepatitis B virus reactivation: Before starting treatment with VOSEVI, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during or after treatment with VOSEVI. This may cause serious liver problems including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B reactivation during and after you stop taking VOSEVI.

ABOUT VOSEVI

VOSEVI is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C (Hep C) infection with or without cirrhosis (compensated) who have:

- genotype 1, 2, 3, 4, 5, or 6 infection and have previously been treated with a Hep C regimen containing an NS5A inhibitor.
- genotype 1a or 3 infection and have previously been treated with a Hep C regimen containing sofosbuvir without an NS5A inhibitor.

Do not take VOSEVI if you take medicines that contain rifampin (Rifater®, Rifamate®, Rimactane®, Rifadin®).

POSSIBLE SIDE EFFECTS OF VOSEVI

VOSEVI can cause serious side effects, including:

- Those in the “Most Important Information about VOSEVI” section.

- **Slow heart rate (bradycardia):** VOSEVI, when taken with amiodarone (Cordarone®, Nexterone®, Pacerone®), a medicine used to treat certain heart problems, may cause serious side effects, including slow heart rate. In some cases slow heart rate has led to death or the need for a pacemaker when amiodarone is taken with medicines containing sofosbuvir. Get medical help right away if you take amiodarone with VOSEVI and get any of the following symptoms:

- fainting or near-fainting
- dizziness or lightheadedness
- not feeling well
- weakness
- extreme tiredness
- shortness of breath
- chest pains
- confusion
- memory problems

The most common side effects of VOSEVI include headache, tiredness, diarrhea, and nausea.

These are not all the possible side effects of VOSEVI. Tell your healthcare provider if you have any new symptoms while taking VOSEVI.

IMPORTANT FACTS

This is only a brief summary of important information about VOSEVI and does not replace talking to your healthcare provider about your condition and your treatment.

BEFORE TAKING VOSEVI

Tell your healthcare provider about all of your medical conditions, including if you:

- have ever had hepatitis B infection
- have liver problems other than hepatitis C infection
- have severe kidney problems or are on dialysis
- are pregnant or breastfeeding, or plan to become pregnant or breastfeed

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider.

VOSEVI and certain medicines may affect each other, or cause side effects.

HOW TO TAKE VOSEVI

- Take VOSEVI with food.
- If you need to take an antacid that contains aluminum or magnesium, take it 4 hours before or 4 hours after you take VOSEVI.

GET MORE INFORMATION

- This is only a brief summary of important information about VOSEVI. Talk to your healthcare provider or pharmacist to learn more.
- Go to VOSEVI.com or call 1-833-4-VOSEVI
- To learn about potential savings, call 1-855-7-MYPATH or go to VOSEVI.com/support-path



VOSEVI, the VOSEVI logo, SUPPORT PATH, GILEAD and the GILEAD logo are trademarks of Gilead Sciences, Inc., or its related companies. All other trademarks referenced herein are the property of their respective owners.

©2018 Gilead Sciences, Inc. All rights reserved. VSVC0025 07/18